



Diversity of the tetracycline resistance gene *tet(M)* and identification of Tn916- and Tn5801-like (Tn6014) transposons in *Staphylococcus aureus* from humans and animals

de Vries, Lisbeth Elvira; Christensen, Henrik; Skov, Robert L.; Aarestrup, Frank Møller; Agersø, Yvonne

Published in:
Journal of Antimicrobial Chemotherapy

DOI:
[10.1093/jac/dkp214](https://doi.org/10.1093/jac/dkp214)

Publication date:
2009

Document version
Early version, also known as pre-print

Citation for published version (APA):
de Vries, L. E., Christensen, H., Skov, R. L., Aarestrup, F. M., & Agersø, Y. (2009). Diversity of the tetracycline resistance gene *tet(M)* and identification of Tn916- and Tn5801-like (Tn6014) transposons in *Staphylococcus aureus* from humans and animals. *Journal of Antimicrobial Chemotherapy*, 64(3), 490-500.
<https://doi.org/10.1093/jac/dkp214>

**Diversity of the Tetracycline Resistance Gene *tet*(M) and
Identification of Tn916- and Tn5801-like (Tn6014) Transposons
in *Staphylococcus aureus* from Human and Animals**

**Lisbeth Elvira de Vries^{1,2}, Henrik Christensen¹, Robert L. Skov³ Frank M.
Aarestrup² and Yvonne Agersø^{2,*}**

¹ Department of Veterinary Disease Biology, University of Copenhagen, Frederiksberg, Denmark

² National Food Institute, The Technical University of Copenhagen, Copenhagen, Denmark

³ Statens Serum Institute, Copenhagen, Denmark

* Corresponding author: Mailing address: National Food Institute, Bülowsvej 27, 1790 Copenhagen
V, Denmark. E-mail: yvoa@food.dtu.dk. Telephone: +4572346273. Fax: +4572346001

Short running title: *tet*(M) in *S. aureus*

Keywords: *tetA*(M), horizontal gene transfer, conjugative transposons, mobile elements

Abstract

Objectives: To analyse the sequence diversity of the tetracycline resistance gene *tet(M)* in *Staphylococcus aureus* from human and animals and to determine mobile elements associated with *tet(M)* in *S. aureus*.

Methods: In total 205 tetracycline resistant isolates were screened for *tet(M)* by PCR. *tet(M)* were sequenced and compared to *tet(M)* deposited with GenBank. Based on phylogenetic analysis isolates were screened for Tn916- and Tn5801-like *xis/int* genes and transposons were confirmed by linking PCR. *spa* typing was performed and selected isolates were used as donors in a filter mating experiment.

Results: Forty-one isolates (21.3 %, 60.7 %, 2.6 % and 4.4 % of the human, pig, poultry and cattle isolates, respectively) were *tet(M)* positive. *tet(M)* was located on Tn5801-like and Tn916-like transposons in humans and on a specific Tn916-like element in animals. Human isolates were of different *spa* types (t034, t008, t037, t051, t065, t078, t318 and t964) corresponding to different clonal complexes (CC398, CC8, CC25 and CC30). Animal isolates were of *spa* type t034, t011 or t0571 corresponding to CC398. *tet(M)* sequence types correlated with CC types. Tn916-like and Tn5801-like (Tn6014) transposons were able to transfer to *S. aureus* recipients.

Conclusion: *S. aureus* of human origin contained diverse *tet(M)* located on Tn916- and Tn5801-like (Tn6014) transposons and *S. aureus* of animal origin contained Tn916-like *tet(M)* genes. This suggest that conjugative transposition play an important role in the evolution and horizontal spread of *tet(M)* in *S. aureus*. This is the first study showing horizontal transfer of Tn5801 (Tn6014).

Introduction

Staphylococcus aureus is part of the normal flora and a frequent cause of infection in humans and many animal species.¹ *S. aureus* are often resistant to tetracycline and two known mechanisms of tetracycline resistance have been identified among staphylococci. Active efflux is a result of acquisition of the genes *tet(K)*, *tet(L)* or *tet(38)*, mainly located on plasmids. Ribosomal protection is conferred by the genes *tet(M)*, *tet(O)*, *tet(S)* or *tet(W)* that are mainly located on different transposons on the chromosome.^{2,3} In addition *tet(U)* has also been found in staphylococci but the mechanism is not fully understood.^{3,4}

tet(M) together with *tet(K)* are the most common genes conferring tetracycline resistance in *S. aureus*.⁵⁻⁸ *tet(M)* is widely distributed among both Gram-positive and Gram-negative bacteria and it has been found in 59 genera.^{3,9} This is probably due to the association of *tet(M)* with integrative and conjugative transposons, facilitating horizontal transfer.¹⁰ Particularly in Gram-positive streptococci and enterococci, *tet(M)* has been found associated with Tn916/1545-like conjugative transposons which form the basis of a family of conjugative transposons that have an extremely broad host range.^{11,12} *tet(M)* associated with Tn916 was originally identified in *Enterococcus faecalis* DS16 and Tn1545 was identified in *Streptococcus pneumoniae*.^{11,13} Recently *tet(M)* was identified on a putative transposon Tn5801 in *S. aureus* Mu50.¹⁴ Tn5801 contains many open reading frames similar to Tn916, but differs by using an integrase (*int*) different from the excisionase/integrase (*xis/int*) present in Tn916 (Figure 1).^{14,15} Other conjugative transposons, like Tn5397 and CW459*tet(M)* have been found to harbour *tet(M)* in *Clostridium difficile* and *Clostridium perfringens*.^{16,17} In addition, *tet(M)* have also been found on different plasmids.²

A limited number of studies concerning the diversity of the *tet(M)* gene have been performed.¹⁸⁻²² In streptococci and enterococci *tet(M)* has been found to be diverse and mainly present on Tn916/Tn1545 like conjugative transposons whereas two different allele types of *tet(M)* from

Lactobacillus has been found to be located mainly on plasmids.¹⁹⁻²² Recently, Agersø *et al.* (2006) showed correlation between diversity of the *tet*(M) DNA sequence and their presence on Tn916, Tn5397 or plasmids in enterococci from different sources in Denmark.²¹

To our knowledge no one has studied the diversity of *tet*(M) and its association with mobile elements in *S. aureus*. A former study found only one out of thirty-four *S. aureus* strains to carry *tet*(M) on Tn916/Tn1545-like transposons.²³ The aim of this study was to analyse sequence diversity of *tet*(M) in *S. aureus* from human and different animals mainly from Denmark, and thereby determine mobile elements associated with *tet*(M) in *S. aureus*.

Material and Methods

Strains

The 205 tetracycline resistant isolates used in this study (Table 1) were identified as *S. aureus* as previously described.²⁴ The 94 human isolates from bacteraemia prospectively collected in Denmark, were selected to represent different phage types and time periods (1957-2002). All human isolates were tested for susceptibility to tetracycline, penicillin, gentamicin, streptomycin, erythromycin and methicillin by tablet diffusion on Danish blood agar as described by the manufacturer (Rosco Neosensitabs, Taastrup). Due to a change in the standard procedure, strains isolated after 1991 were additionally tested for susceptibility to fusidic acid, ciprofloxacin and rifampicin in the same way (Table 2).

Of the 111 animal isolates (Table 1), 39 poultry²⁵, 27 pig and 2 lamb isolates were diagnostic submissions to either The National Veterinary Institute or to The National Food Institute, Technical University of Denmark. One pig isolate (9b) was obtained from a healthy pig in 2007. All animal isolates were tested for susceptibility to tetracycline, penicillin, streptomycin, erythromycin, ciprofloxacin, spectinomycin, tiamulin, trimethoprim, ceftiofur, chloramphenicol, florfenicol and

93 sulphamethoxazole by use of sensititre method as described previously (Table 2).²⁵ One pig isolate
94 9b was confirmed to be MRSA by *mecA* PCR.²⁶

95 **Screening and sequencing of *tet(M)***

96 All 205 isolates were screened for *tet(M)* by PCR as described previously (Table 1).²⁷ For all 41
97 *tet(M)* positive isolates three or four overlapping PCR fragments covering *tet(M)* including a
98 downstream region were amplified and used as templates for sequencing. Different sequencing
99 strategies were used in parallel as outlined in Figure 2. DNA Taq polymerase (Ampliqon, Denmark)
100 was used for all PCR amplifications. Sequencing was performed by Macrogen, Korea.²⁸ Primers
101 526, 540, 324, 525, 266, 323, 307, 323, 307 and 1756 were used (Table 3).

102 **Detection of Tn916-like and Tn5801-like transposons**

103 The presence of the *int* genes specific for Tn5801 was detected by PCR using primers 1811 and
104 1812 (product size 722 bp) and DNA Taq polymerase from Ampliqon. Tn916-like *xis* genes were
105 detected by PCR as previously described.²⁹ Tn916-*xis* screening PCR products from human isolates
106 8797 and 5377 were sequenced using amplification primers (327-328) and the Tn5801-*int* screening
107 product from human isolate 1680 was sequenced with amplification primers (1811-1812). Primers
108 are listed in Table 3.

109 Long PCR linking *tet(M)* to Tn916-like *xis* or Tn5801-like *int* was performed with Phusion™ High-
110 Fidelity DNA Polymerase (Finnzymes, Finland) using condition as recommended by the
111 manufacturer. Primers 328 and 804 (Table 3) were used for long PCRs linking *tet(M)* with Tn916-
112 like *xis*. PCR conditions were 30 sec at 98°C followed by 30 cycles of 10 sec at 98°C, 30 sec at
113 51°C, 85 sec at 72°C and a final extension for 10 min at 72°C. Primers 709 and 1812 (Table 3) were
114 used for long PCR linking Tn5801-like *int* with *tet(M)* using conditions of initial denaturation for
115 30 sec at 98°C followed by 30 cycles of 10 sec at 98°C, 30 sec at 61°C, 144 sec at 72°C and a final

extension for 10 min at 72 °C. The *tet(M)*-Tn5801-like *int* product from of human isolate 1680 was sequenced with primers 1812 and 1835-1840 (GenBank submission no. EU918655).

In all PCRs the reference strains *E. faecalis* DS16³⁰ and *S. aureus* Mu50¹⁴ were used as positive control for the presence of Tn916- and Tn5801- like transposons, respectively. As negative control the reference strain containing the other transposon was used.

Phylogenetic analysis

GenBank was searched for full length *tet(M)* genes based on the definition that *tet(M)* genes share ≥80% similarity on the amino acid level.² Fifty two unique gene sequences were selected to represent different species from different hosts. A Neighbor Joining (NJ) tree based on a multiple alignment of 41 sequences obtained in this study and 52 *tet(M)* genes from GenBank (1920 bp) was constructed in Clustal X³¹ and visualized by MEGA 3.1.³² The tree was rooted with the *tet(O)* gene (GenBank/EMBL/DDBJ accession nr. Y07780) as outgroup. Another tree based on the 450 bp region downstream of *tet(M)* was constructed in the same way. Sequences were compared pairwise with the EMBOSS program water used for local alignments.^{33 34}

spa and MLST typing

All *tet(M)* positive isolates were *spa* typed (Table 2) using primers and conditions recommended by SeqNet.³⁵ The *spa* types were determined using BioNumerics 4.61 (Applied maths, Sint-Martens-Latem, Belgium). Two human and one pig isolate (1591, 34801 and 9b) were also MLST typed as recommended by MLSTnet.³⁶

Clustering of *tet(M)* verses *spa* types in clonal complexes

CC types were deduced from the *spa* types by using information available from the Ridom *Spa* Server and from MLSTnet.^{37 38} In cases where the CC type could not be deduced from the *spa* type

(isolate 1591, 34801 and 9b), the CC type was determined from the MLST type by using the eBURSTv3 algorithm.³⁹ Related *spa* types within different CC types were revealed by using the Minimum Spanning Tree method in BioNumerics, cut off distance ≤ 3 .

Filter mating

Filter mating experiments were performed as described previously⁴⁰ using nine human isolates (1591, 1680, 21995, 34148, 34168, 34801, 35366, 4520, 8797) and five animal isolates (7413532-2, 7611472-1, 9877324-3, USA42, 9b) as donors and the two *S. aureus* recipients, R1 (8794RF)⁴¹ and R2 (RN4220RF).⁴² The detection limit of transconjugants and the rates of spontaneous mutations were calculated for each of the mating experiment. In all experiments the donors tend to grow faster than the recipient, therefore the transfer rates and the detection limits were calculated as transconjugants per recipient. Transconjugants were selected on brain heart infusion agar plates (Becton, Dickinson and Company, USA), supplemented with 8 mg/L of tetracycline, 12.5 mg/L of rifampicin and 12.5 mg/L of fusidic acid. The numbers of donors and recipient were counted on brain heart infusion agar plates supplemented with 8 mg/L of tetracycline or 12.5 mg/L of rifampicin and 12.5 mg/L of fusidic acid, respectively. Transconjugants were further verified by *spa* typing and screened for *tet*(M) by PCR. Long PCR linking *tet*(M) to Tn916-like *xis* or Tn5801-like *int* verified that *tet*(M) was present on either Tn916- or Tn5801-like transposons in the transconjugants.

Results

Screening *S. aureus* isolates for *tet*(M)

Out of 205 tetracycline resistant *S. aureus* isolates, 20 human and 21 animal isolates were shown to be positive for *tet*(M) by PCR (Table 1). The highest prevalence of *tet*(M) was found among

isolates from pigs with 60.7 % *tet*(M) positives compared to 21.3% in humans, 4.3% in cattle and 2.6% in poultry.

Sequencing of *tet*(M)

The *tet*(M) gene including a downstream region from 41 *S. aureus* isolates were sequenced according to the strategies shown in Figure 2. Comparing all 41 *tet*(M) gene sequences (1920 bp) revealed 6 unique sequence types, of which one type was sequenced with strategy 1 and the other five types were sequenced with either strategy 2a or 2b.

Phylogenetic analysis predicts mobile elements associated with *tet*(M)

The result of the phylogenetic analysis is shown in Figure 3. The sequences fell into three groups. All staphylococci sequences including the 41 *tet*(M) from *S. aureus*, Tn5801 *tet*(M) from *S. aureus* Mu50 (BA000017) and Tn916 *tet*(M) from *E. faecalis* DS16 (U09422) fell into group II.²¹ Tn5397 *tet*(M) from *C. difficile* (AF333235) and two similar Tn5397-like *tet*(M) from *E. faecium* were contained in group I and Tn1545 *tet*(M) from *E. faecalis* (X04388) and other composite transposons (Tn2009 and Tn5251) fell into group III.

Based on similarity, the 41 *tet*(M) genes from *S. aureus* (consisting of 6 sequence types) were divided into three subgroups within group II (Figure 3). Subgroups 1 and 2 were identical or highly related (98.8-100% similarity at DNA level) to *tet*(M)-Tn916 from *E. faecalis* DS16, however subgroup 2 formed an individual branch supported with a bootstrap of 100%. Sequences of subgroup 3 were identical to *tet*(M) Tn5801 from *S. aureus* Mu50 (BA000017). This indicates that *tet*(M) of subgroups 1 and 2 were located on Tn916-like transposons and that *tet*(M) sequences of subgroup 3 were located on the putative transposon Tn5801. This was further supported by a phylogenetic tree based on the 450 bp region downstream of *tet*(M) that divided the sequences into

two groups (data not shown). One group was identical or highly related to the downstream region of Tn916 (99.8-100%) and the second group was identical to the downstream region of Tn5801.

***S. aureus* tet(M) genes are located on Tn916-like and Tn5801-like transposons**

The presence of *tet*(M) on Tn916-like transposons in subgroups 1 and 2 and on the putative transposon Tn5801 in subgroup 3 (Figure 3) was confirmed by PCR. All isolates from subgroup 1 and 2 were positive for Tn916-*xis* and negative for Tn5801-*int* and all isolates from subgroup 3 were positive for Tn5801-*int* and negative for Tn916-*xis* (data not shown). Two *xis* and one *int* PCR screening products from the human isolates 8797, 5377 and 1680 representing subgroup 1, 2 and 3 respectively were sequenced. Both *xis* sequences were 100% identical to the corresponding *xis* sequence from Tn916 in *E. faecalis* DS16 and the *int* sequence were 100% identical to the corresponding *int* sequence from Tn5801 in *S. aureus* Mu50.

For all isolates with *tet*(M), linking PCR confirmed that the Tn916-*xis* and Tn5801-*int* genes detected in the PCR screen were actually located in the same element as *tet*(M) (Figure 4A). The DNA sequence of *tet*(M)-*int* (GenBank accession no. EU918655) from human isolate 1680 (subgroup 3) had 99.9% similarity with the corresponding sequence in Tn5801 (BA000017). Thus *tet*(M) of subgroups 1 and 2 are located on Tn916-like elements and *tet*(M) of subgroup 3 is located on Tn5801-like elements.

Dissemination of *tet*(M) within *S. aureus* of human and animal origin

In Table 2, *spa* types and corresponding CC types are shown. Most animal isolates had *spa* type t034 except two pig isolates of *spa* type t011 and one pig isolate with *spa* type t571, all belonging to CC398.⁴³ The human isolates had different *spa* types: t034 (CC398), t008, t037 and t051 (CC8), t065 (CC45), t078 (CC25), t318 and t964 (CC30) and t668 (CC5). Thus isolates of *spa* type t034 (CC398) were found both in different animals and in humans.

In order to compare how the different *tet(M)* genes may have been disseminated within and between different CC types of *S. aureus*, the *tet(M)* sequences were grouped according to their CC type (Figure 5). Figure 5 shows a clear correlation between different *tet(M)* sequence types and different CC types of *S. aureus*. Moreover, *tet(M)* of subgroup 3 was identical to *tet(M)*-Tn5801 from *S. aureus* Mu50 belonging to CC5, indicating horizontal transfer of Tn5801 between CC5 and CC8 (Figure 5C).

Horizontal transfer of Tn916- and Tn5801-like (Tn6014) transposons

To test whether the identified Tn916-like and Tn5801-like transposons were functional conjugative transposons, filter mating experiments with 14 selected isolates as donors and two *S. aureus* recipients were performed. The detection limits for the mating experiments were between 2.7×10^{-10} to 1.5×10^{-9} transconjugants/recipient except for mating with donor 1591 and recipient R1 (8794) where the detection limit was 2.5×10^{-8} transconjugants/recipient. Spontaneous mutations to rifampicillin and fudisic acid were observed only for donor 8797 ($1.4\text{--}1.9 \times 10^{-8}$) and donor 34801 ($0.9\text{--}8.5 \times 10^{-10}$). Transconjugants conferring resistance to tetracycline, but not containing *tet(M)* were observed only in matings with donor 9877324-3 to both recipients (1×10^{-7} and 9×10^{-9} transconjugants/recipient, respectively).

Tn916-like *tet(M)* from the human isolates 34801 and 35366 from subgroup 1 were able to transfer to R1 (8794RF) at transfer rates of 1×10^{-9} transconjugants/recipient and 3×10^{-8} transconjugants/recipient, respectively. Transfer from isolate 35366 into R2 (RN4220RF) was also observed (1×10^{-9} transconjugants/recipient). Tn5801-like *tet(M)* of the human isolate 1680 from subgroup 3 was able to transfer to R2 (RN4220RF) with a transfer rate of 1×10^{-9} transconjugants/recipient (Figure 4B and 4C). This transfer was in addition to *spa* typing verified by a PFGE analysis showing the recipient R2 and the transconjugant (1680R2_4) to have the same PFGE pattern distinct of the donor (1680) after *Sma*I digestion (data not shown). The Tn5801-like

element from human isolate 1680 was therefore registered as a novel conjugative transposon, Tn6014 in the Transposon Nomenclature Database from the UCL Eastman Dental Institute, London (<http://www.ucl.ac.uk/eastman/tn/>).⁴⁴

Discussion

The screening of tetracycline resistant *S. aureus* isolates showed the highest prevalence of *tet*(M) among the tetracycline resistant pig isolates (60.7 %) and the lowest in tetracycline resistant poultry (2.6 %) and bovine-mastitis (4.4 %) isolates. All animal isolates belonged to CC398 that has recently emerged as a Methicillin-resistant clone in the Netherlands and other countries including Denmark.⁴⁵⁻⁴⁷ Beside the animal isolates four of the twenty one human isolates also belonged to CC398 and contained the same Tn916-like *tet*(M) gene as all the animal isolates. CC398 isolates are usually tetracycline resistant and a recent study detected *tet*(M) in all methicillin-resistant *S. aureus* (MRSA) CC398 studied from human and companion animals in Germany and Austria.^{43, 45, 47} This suggests that a Tn916-like *tet*(M) was integrated and adapted early in the evolution of the clone and may be disseminated vertically within CC398.

In our study one of the human CC398 isolates dates back to 1992. The two CC398 isolates from cattle are from the beginning of the 1990s and the turkey CC398 isolates is from 1998. The rest of CC398 were isolated between 2000 and 2008. Thus already in the early 1990s inter-species transmission of CC398 may have occurred. Whether the high occurrence of CC398 among the pig isolates found in this study reflects that pigs are the main reservoir for CC398 *tet*(M) is unknown.

Sequence analysis divided the sequenced *tet*(M) into three subgroups corresponding to two different transposons. *tet*(M) of subgroup 1 and 2 were located on Tn916-like transposons and subgroup 3 was located on Tn5801-like transposons. Subgroup 1 contained human isolates from different *spa* types (and phage types) and all animal isolates from the time span 1959-2007. The sequence variations of *tet*(M) within subgroup 1 correlated with different CC types of *S. aureus* (see

Figure 5A) which supports the general idea that *S. aureus* of different lineages are not very good at sharing DNA. Subgroup 2 formed an individual branch with five identical *tet*(M) sequences, all from human isolates of phage type 94/96 isolated between 1970-1992. All were shown to have the same *spa* type (t078) belonging to CC25. Thus *tet*(M) of subgroup 2 appears to have been integrated in this clone over 30 years ago without changing. PCR mapping of the elements in subgroup 2 were of expected size (data not shown), indicating that all the ORFs necessary to conjugate were present. Whether or not this new Tn916-like transposon is functional is however not clear. Subgroup 3 consisted of seven isolates from 1957-2000 with identical *tet*(M), different phage types and *spa* types all belonging to CC8. Comparing the isolates within this group shows a correlation between *spa* type and resistance pattern (see Table 2). The three isolates from *spa* type t037 (all from 2000) were resistant to the same seven antimicrobial agents including Methicillin and were suspected to be from the same outbreak. The two other *spa* types in this group, t008 and t051 (1957-1963) are clonally related and are only resistant to 3 and 5 agents respectively, the latter was Methicillin-resistant. These differences may be time dependent or reflect resistance profiles in different sub-clones of CC8.

As shown by the phylogenetic tree in Figure 3, *S. aureus tet*(M) sequences belonging to subgroups 1A, 1C and 3 were identical to *tet*(M) of *Streptococcus agalactiae* (AAJQ1000009), Tn916 -*tet*(M) from *E. faecalis* DS16 (U09422) and to *tet*(M) found in *Streptococcus agalactiae* COH1 (NZ_AAJR01000021), respectively. This indicates horizontal transfer of *tet*(M) from subgroup 1 and subgroup 3 between *S. aureus* and other Gram-positive species of enterococci and streptococci. Previously horizontal transfer of Tn916-like *tet*(M) from *Bacillus cereus* group into *S. aureus* has been shown.²⁹ Clustering of CC *spa* types versus *tet*(M) sequences type suggested that horizontal transfer of Tn5801 between different CC types of *S. aureus* has occurred.

The filter mating experiment showed that the Tn916-like element from subgroup 1A (CC398) could be transferred into both recipient strains (8794RF, CC121 and RN4220RF, CC8) whereas the Tn916-like element from subgroup 1D (CC5) was only transferred into one of the recipient strains (8794RF, CC121). The Tn5801-like element, Tn6014 from subgroup 3 (CC8) was transferred into the other recipient strain (RN4220RF, CC8). The new *tet(M)*-like elements of subgroup 2 did not transfer into any of the recipients. Although, both recipients are known to be very good in taking up foreign DNA, transconjugants were obtained with very low frequencies. Recently the restriction-modification system, *SauI* was suggested to control horizontal gene transfer between *S. aureus* of different lineages.⁴⁸ RN4220RF was shown to have a mutation in this system making it able to take up DNA from different CC types. However, in our study RN4220RF only received *tet(M)* from two of the 14 tested donors which indicates that other factors may also play a role or that transfer occurs at rates below our detection limit.

Staphylococci tet(M) were only located in part of the phylogenetic tree associated with the well characterized conjugative transposon Tn916 and with Tn5801 described in *S. aureus* Mu50 and Mu3^{14, 49} (group II, Figure 3). *tet(M)* from other Gram-positive bacteria were distributed in the whole tree and were besides Tn916-like elements associated with Tn5397 (group I) and/or composite transposons like Tn1545, Tn2009 and/or Tn5251 (group III). Moreover, *tet(M)* from *E. faecium* (DQ223243 and DQ223244) has also been found on plasmids (group I).²¹ Thus *tet(M)* from *S. aureus* appear to be less diverse than *tet(M)* from other Gram-positive bacteria.

The predicted mobile elements associated with *tet(M)* in *S. aureus* from different origins were confirmed experimentally by long PCR. The same approach was also used successfully in a previous study of the diversity of *tet(M)* among Enterococci.²¹ Thus in general *tet(M)* appear to be more related to its mobile element than to the host species, however module exchange between transposons and recombination within *tet(M)* may also have occurred.^{10, 21} Module exchange seems

to be the case for putative transposon CW459*tet*(M) from *Clostridium peringens*.¹⁶ In this transposon *tet*(M) is highly related to Tn916-*tet*(M) whereas the rest of the transposon sequence is more related to Tn5801.¹⁵

In conclusion, we have used the diversity of the *tet*(M) gene to determine associated mobile elements in *S. aureus* from human and different animal origin. *S. aureus* of human origin was shown to contain diverse *tet*(M) genes located on Tn916-like and Tn5801-like conjugative transposons that corresponded with different CC types. *S. aureus* of different animal origin contained one specific type of *tet*(M) located on Tn916-like elements, all belonging to CC type 398. This is the first report showing that a Tn5801-like element, Tn6014 can transfer between *S. aureus* isolates.

Acknowledgements

We want to thank Maria Louise Johannsen, Jacob Dyring Jensen and Hanne Mordhorst for excellent technical assistance. Part of this work was presented as a poster at the 47th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC), 2007 and at the 1st International ASM meeting on Antimicrobial Resistance in Zoonotic bacteria and foodborne pathogens, 2008.

Funding

This study was founded by a grant from the The Danish Research Council for Technology and Production Sciences (274-05-0117).

Transparency declarations

None to declare

Reference

1. Aarestrup FM, Schwarz S. Antimicrobial Resistance in Staphylococci and Streptococci of Animal Origin. In: Aarestrup FM, ed. *Antimicrobial Resistance in Bacterial of Animal Origin*. Washington: ASM, 2006; 187-212.
2. Chopra I, Roberts M. Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiol Mol Biol Rev* 2001; **65**: 232-60.
3. Roberts MC. Update on acquired tetracycline resistance genes. *FEMS Microbiol Lett* 2005; **245**: 195-203.
4. Ridenhour MB, Fletcher HM, Mortensen JE, et al. A novel tetracycline-resistant determinant, *tet(U)*, is encoded on the plasmid pKq10 in *Enterococcus faecium*. *Plasmid* 1996; **35**: 71-80.
5. Bismuth R, Zilhao R, Sakamoto H, et al. Gene heterogeneity for tetracycline resistance in *Staphylococcus* spp. *Antimicrob Agents Chemother* 1990; **34**: 1611-4.
6. Schmitz FJ, Krey A, Sadurski R, et al. Resistance to tetracycline and distribution of tetracycline resistance genes in European *Staphylococcus aureus* isolates. *J Antimicrob Chemother* 2001; **47**: 239-40.
7. Warsa UC, Nonoyama M, Ida T, et al. Detection of *tet(K)* and *tet(M)* in *Staphylococcus aureus* of Asian countries by the polymerase chain reaction. *J Antibiot (Tokyo)* 1996; **49**: 1127-32.
8. Schwarz S, Roberts MC, Werckenthin C, et al. Tetracycline resistance in *Staphylococcus* spp. from domestic animals. *Vet Microbiol* 1998; **63**: 217-27.
9. Distribution of *tet(M)* in Gram-positive and Gram-negative bacteria.
<http://faculty.washington.edu/marilynr/> (23 September 2008, date last accessed).
10. Burrus V, Pavlovic G, Decaris B, et al. Conjugative transposons: the tip of the iceberg. *Mol Microbiol* 2002; **46**: 601-10.
11. Rice LB. Tn916 family conjugative transposons and dissemination of antimicrobial resistance determinants. *Antimicrob Agents Chemother* 1998; **42**: 1871-7.
12. Clewell DB, Francia MV. Conjugation in Gram-positive bacteria. In: Funnell BE, Phillips G.J.E., eds. *Plasmid Biology*. Washington DC: American Society for Microbiology, 2004; 227-56.
13. Courvalin P, Carrier C. Tn1545: a conjugative shuttle transposon. *Mol Gen Genet* 1987; **206**: 259-64.
14. Kuroda M, Ohta T, Uchiyama I, et al. Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet* 2001; **357**: 1225-40.

- 355 15. Ito T, Okuma K, Ma XX, et al. Insights on antibiotic resistance of *Staphylococcus aureus*
356 from its whole genome: genomic island SCC. *Drug Resist Updat* 2003; **6**: 41-52.
- 357 16. Roberts AP, Johanesen PA, Lyras D, et al. Comparison of Tn5397 from *Clostridium difficile*,
358 Tn916 from *Enterococcus faecalis* and the CW459tet(M) element from *Clostridium*
359 *perfringens* shows that they have similar conjugation regions but different insertion
360 and excision modules. *Microbiology* 2001; **147**: 1243-51.
- 361 17. Lyras D, Rood JJ. Genetic organization and distribution of tetracycline resistance determinants
362 in *Clostridium perfringens*. *Antimicrob Agents Chemother* 1996; **40**: 2500-4.
- 363 18. Oggioni MR, Dowson CG, Smith JM, et al. The tetracycline resistance gene *tet*(M) exhibits
364 mosaic structure. *Plasmid* 1996; **35**: 156-63.
- 365 19. Doherty N, Trzcinski K, Pickerill P, et al. Genetic diversity of the *tet*(M) gene in tetracycline-
366 resistant clonal lineages of *Streptococcus pneumoniae*. *Antimicrob Agents Chemother*
367 2000; **44**: 2979-84.
- 368 20. Huys G, D'Haene K, Collard JM, et al. Prevalence and molecular characterization of
369 tetracycline resistance in *Enterococcus* isolates from food. *Appl Environ Microbiol*
370 2004; **70**: 1555-62.
- 371 21. Agersø Y, Pedersen AG, Aarestrup FM. Identification of Tn5397-like and Tn916-like
372 transposons and diversity of the tetracycline resistance gene *tet*(M) in enterococci from
373 humans, pigs and poultry. *J Antimicrob Chemother* 2006; **57**: 832-9.
- 374 22. Gevers D, Danielsen M, Huys G, et al. Molecular characterization of *tet*(M) genes in
375 *Lactobacillus* isolates from different types of fermented dry sausage. *Appl Environ*
376 *Microbiol* 2003; **69**: 1270-5.
- 377 23. Poyart-Salmeron C, Trieu-Cuot P, Carlier C, et al. Nucleotide sequences specific for Tn1545-
378 like conjugative transposons in pneumococci and staphylococci resistant to
379 tetracycline. *Antimicrob Agents Chemother* 1991; **35**: 1657-60.
- 380 24. Aarestrup FM, Wegener HC, Rosdahl VT, et al. Staphylococcal and other bacterial species
381 associated with intramammary infections in Danish dairy herds. *Acta Vet Scand* 1995;
382 **36**: 475-87.
- 383 25. Aarestrup FM, Agersø Y, Ahrens P, et al. Antimicrobial susceptibility and presence of
384 resistance genes in staphylococci from poultry. *Vet Microbiol* 2000; **74**: 353-64.
- 385 26. Multiplex PCR for the detection of the *mecA* gene. <http://www.crl-ar.eu/data/images/meca->
386 [pcr_protocol%2006.02.08.pdf](http://www.crl-ar.eu/data/images/meca-) (24 September 2008, date last accessed).
- 387 27. Aarestrup FM, Agersø Y, Gerner-Smidt P, et al. Comparison of antimicrobial resistance
388 phenotypes and resistance genes in *Enterococcus faecalis* and *Enterococcus faecium*
389 from humans in the community, broilers, and pigs in Denmark. *Diagn Microbiol Infect*
390 *Dis* 2000; **37**: 127-37.

- 391 28. Macrogen. http://www.macrogen.com/eng/sequencing/sequence_main.jsp (23 September
392 2008, date last accessed).
- 393 29. Agersø Y, Jensen LB, Givskov M, et al. The identification of a tetracycline resistance gene
394 *tet(M)*, on a Tn916-like transposon, in the *Bacillus cereus* group. *FEMS Microbiol Lett*
395 2002; **214**: 251-6.
- 396 30. Clewell DB. Movable genetic elements and antibiotic resistance in enterococci. *Eur J Clin*
397 *Microbiol Infect Dis* 1990; **9**: 90-102.
- 398 31. Thompson JD, Gibson TJ, Plewniak F, et al. The CLUSTAL_X windows interface: flexible
399 strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic*
400 *Acids Res* 1997; **25**: 4876-82.
- 401 32. Kumar S, Tamura K, Nei M. MEGA3: Integrated software for Molecular Evolutionary
402 Genetics Analysis and sequence alignment. *Brief Bioinform* 2004; **5**: 150-63.
- 403 33. Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open Software
404 Suite. *Trends Genet* 2000; **16**: 276-7.
- 405 34. EMBOSS Pairwise Alignment Algorithms. <http://www.ebi.ac.uk/emboss/align/> (23 September
406 2008, date last accessed).
- 407 35. *spa* typing. http://www.seqnet.org/pdf/Ridom_spa_sequencing.pdf (23 September 2008, date
408 last accessed).
- 409 36. MLST typing. <http://saureus.mlst.net/misc/info.asp#experimental> (23 September 2008, date
410 last accessed).
- 411 37. Ridom SpaServer. <http://spaserver2.ridom.de/mlst.shtml> (23 September 2008, date last
412 accessed).
- 413 38. *Staphylococcus aureus* MLST database. <http://saureus.mlst.net/> (23 September 2008, date last
414 accessed).
- 415 39. eBURST. <http://eburst.mlst.net/> (28 September 2008, date last accessed).
- 416 40. Hammerum AM, Jensen LB, Aarestrup FM. Detection of the *satA* gene and transferability of
417 virginiamycin resistance in *Enterococcus faecium* from food-animals. *FEMS*
418 *Microbiol Lett* 1998; **168**: 145-51.
- 419 41. Stobberingh EEWK. Restriction-deficient mutants of *Staphylococcus aureus*. *J Gen*
420 *Microbiol* 1977; **99**: 359-67.
- 421 42. Kreiswirth BN, Lofdahl S, Betley MJ, et al. The toxic shock syndrome exotoxin structural
422 gene is not detectably transmitted by a prophage. *Nature* 1983; **305**: 709-12.
- 423 43. Wulf MW, Sørum M, van NA, et al. Prevalence of methicillin-resistant *Staphylococcus*
424 *aureus* among veterinarians: an international study. *Clin Microbiol Infect* 2008; **14**:
425 29-34.

- 426 44. Roberts AP, Chandler M, Courvalin P, et al. Revised nomenclature for transposable genetic
427 elements. *Plasmid* 2008; **60**: 167-73.
- 428 45. Witte W, Strommenger B, Stanek C, et al. Methicillin-resistant *Staphylococcus aureus* ST398
429 in humans and animals, Central Europe. *Emerg Infect Dis* 2007; **13**: 255-8.
- 430 46. Huijsdens XW, van Dijke BJ, Spalburg E, et al. Community-acquired MRSA and pig-
431 farming. *Ann Clin Microbiol Antimicrob* 2006; **5**: 26.
- 432 47. Guardabassi L, Stegger M, Skov R. Retrospective detection of methicillin resistant and
433 susceptible *Staphylococcus aureus* ST398 in Danish slaughter pigs. *Vet Microbiol*
434 2007; **122**: 384-6.
- 435 48. Waldron DE, Lindsay JA. Sau1: a novel lineage-specific type I restriction-modification
436 system that blocks horizontal gene transfer into *Staphylococcus aureus* and between *S.*
437 *aureus* isolates of different lineages. *J Bacteriol* 2006; **188**: 5578-85.
- 438 49. Hiramatsu K, Aritaka N, Hanaki H, et al. Dissemination in Japanese hospitals of strains of
439 *Staphylococcus aureus* heterogeneously resistant to vancomycin. *Lancet* 1997; **350**:
440 1670-3.
- 441 50. Clewell DB, Flannagan SE, Jaworski DD. Unconstrained bacterial promiscuity: the Tn916-
442 Tn1545 family of conjugative transposons. *Trends Microbiol* 1995; **3**: 229-36.
- 443 51. Vintov J, Aarestrup FM, Zinn CE, et al. Association between phage types and antimicrobial
444 resistance among bovine *Staphylococcus aureus* from 10 countries. *Vet Microbiol*
445 2003; **95**: 133-47.

Legends to Figures

Figure 1: Comparison of the 18 kb conjugative transposon Tn916 (U09422) and the 25 kb putative transposon Tn5801 (BA000017/NC002758). Besides *tet(M)* (dark grey arrows) they both consist of 3 structural domains containing genes associated with conjugation, regulation (light grey arrows) and excision/integration (dotted arrows).^{11, 15, 50} Both elements contain an integrase (*int*) gene, however these *int* genes are very different. In addition, Tn916 also contains an annotated excisionase (*xis*) gene (black arrow). Tn5801 contains several open reading frames (white arrows) whose functions are unknown. *sav415* may however encode a transposase.¹⁵ The relation between open reading frames of Tn916 and Tn5801 are shown in percent identity on nucleotide level calculated with the EMBOSS program water.³⁴

Figure 2.: Amplification and sequencing strategy for *tet(M)*. Top: The *tet(M)* gene sequence including a downstream region. Primers used for amplification and sequencing are illustrated with arrows. Bottom: Two different sequencing strategies using different combination of primers. Strategy 1: Sequences from 7 human isolates (213, 229, 1680, 1742, 33597, 34148, 34168) were obtained. Strategy 2a: 3 human, 8 pig, 1 lamb and 1 bovine isolate (22034, 35414, 35679, 9b, 7215311, 7311242, 7413093-4, 7413714-1, 7512986-1, 7611472.1, 7611995-1, 7612628-4, sw356). Strategy 2b: 10 human, 9 pig, 1 turkey and 1 bovine isolate (617, 1591, 4520, 4865, 5331, 5377, 8797, 21995, 35801, 35366, 7215190-1, 7312330-1, 7412791, 7413532-2, 7413727, 74140355-2, 7512166-1, 7611280, 7711730-1, 9877324-3, usa42).

Figur 3: Phylogenetic gene tree of *tet(M)*. Bootstrap values are indicated at branch points (out of 1000 generated NJ trees). Group I is supported by a bootstrap value of 87.8%, group II by a

bootstrap value of 48.7% at the first branching and 93.7% at the second branching and group 3 by a bootstrap value of 99.4%.

Figure 4: PCR products linking *tet*(M) to Tn916-like *xis* and Tn5801-like *int* genes with the expected size of 2835 bp and 4820 bp respectively. **A:** A representative from every *tet*(M) type from different origins are shown. Lane 1: 9877324-3 (turkey), 2: USA42 (cattle), 3: 7611472-1 (lamb), 4: 9b (pig), 5: 21995 (human), 6: 617 (human), 7: 8797 (human), 8: 34801 (human), 9: 4520 (human), 10: positive control (*E. faecalis* DS16), 11: negative control (*S. aureus* Mu50), 12: 1680 (human), 13: positive control (*S. aureus* Mu50), 14: negative control (*E. faecalis* DS16). M: Gene Ruler 1Kb ladder from Fermentas. **B:** Transconjugants (TC), donors (D) and recipients showing horizontal transfer of Tn916-like transposons. Lane 1: 34801_1_R1 (TC), 2: 34801 (D), 3: 35366_1_R1 (TC), 4: 35366_1_R2, 5: 35366 (D), 6: R1 (8794RF), 7: R2 (RN4220RF), 8: positive control (*E. faecalis* DS16), 9: negative control (*S. aureus* Mu50). **C:** TC, D and recipient showing horizontal transfer of the Tn5801-like transposon Tn6014. Lane 1: 1680R2_4 (TC), 2: 1680 (D), 3: R2 (RN4220RF), 4: positive control (*S. aureus* Mu50), 5: negative control (*E. faecalis* DS16).

Figure 5: CC *spa* type clustering versus different *tet*(M) sequence types in *S. aureus*. Predicted horizontal gene transfer of *tet*(M) is illustrated by an arrow. Related *spa* types are illustrated with a black line: *spa* type t034 is related to t011 by a deletion of two repeats in t011 compared to t034. Furthermore, t034 is related to t571 by a deletion of one repeat in t571 compared to t034. *spa* type t051 is related to t008 by a deletion of one repeat in t008 compared to t501. *spa* types t964 and t318 shared 8 out of their 9 or 10 repeats, respectively and one repeat varies by one substitution.

Table 1: Origin, source, phage types, country, year and numbers of *S. aureus* isolates screened for *tet(M)* by PCR and the number found positive for *tet(M)*.

<i>Source</i>	<i>Human^a</i>	<i>Pig</i>	<i>Poultry</i>	<i>Sheep</i>	<i>Cattle^b</i>
<i>Phage types</i>	80 complex, gr1, gr2, gr3, 83A, 94/96, 95, MIX, NI	ND	ND	ND	ND
<i>Country/Region</i>	Denmark	Denmark	Denmark	Denmark	Europe & US
<i>Year</i>	1957-2002	2000-2007	1994-1998	2004-2005	Early 1990s
<i>Isolates</i>	94	28	39	2	42 ⁵¹
<i>tet(M) positive</i>	20 (21.3%)	17 (60.7%)	1 (2.6%)	1 (50%)	2 (4.3%)

a: bacteraemia, b: mastitis, ND: Not determined

525 **Table 2.** *S. aureus* strains with *tet*(M) used in this study

Strain	Source/year	Other resistance phenotypes	Mobile element carrying <i>tet</i> (M)	<i>spa</i> type/CC types	GenBank accession number
213	Human/1957	Pen ^R , Str ^R	Tn5801-like	t008/CC8	EU918651
229	Human/1957	Pen ^R , Str ^R	Tn5801-like	t008/CC8	EU918652
1680	Human/1963	Pen ^R , Str ^R , Ery ^R , Met ^R	Tn5801-like (Tn6014)	t051/CC8	EU918655
1742	Human/1963	Pen ^R , Str ^R	Tn5801-like	t008/CC8	EU918656
33597 ^b	Human/2000	Pen ^R , Str ^R , Gen ^R , Ery ^R , Met ^R , Cip ^R	Tn5801-like	t037/CC8	EU918664
34148 ^b	Human/2000	Pen ^R , Str ^R , Gen ^R , Ery ^R , Met ^R , Cip ^R	Tn5801-like	t037/CC8	EU918665
34168 ^b	Human/2000	Pen ^R , Str ^R , Gen ^R , Ery ^R , Met ^R , Cip ^R	Tn5801-like	t037/CC8	EU918666
4520	Human/1970	Pen ^R	Tn916-like	t078/CC25	EU918657
4865	Human/1970	Pen ^R	Tn916-like	t078/CC25	EU918658
5331	Human/1971	Pen ^R	Tn916-like	t078/CC25	EU918659
5377	Human/1971	Pen ^R	Tn916-like	t078/CC25	EU918660
22034	Human/1992	Pen ^R , Ery ^R , Cip ^R	Tn916-like	t078/CC25	EU918663
34801	Human/2001	Pen ^R , Fus ^R	Tn916-like	t688/CC5 ^a	EU918667
8797	Human/1977	Pen ^R	Tn916-like	t065/CC45	EU918661
617	Human/1959	Pen ^R	Tn916-like	t318/CC30	EU918653
1591	Human/1962	Pen ^R	Tn916-like	t964/CC30 ^a	EU918654
21995	Human/1992	Pen ^R	Tn916-like	t034/CC398	EU918662
35366	Human/2001	Pen ^R	Tn916-like	t034/CC398	EU918668
35414	Human/2001		Tn916-like	t034/CC398	EU918669
35679	Human/2001	Pen ^R	Tn916-like	t034/CC398	EU918670
9877324-3	Turkey/1998	Pen ^R	Tn916-like	t034/CC398	EU918671
USA42	Cattle	Pen ^R , Str ^R , Ery ^R , Chl ^R , Sul ^R	Tn916-like	t034/CC398	EU918673
Sw356	Cattle	Str ^R , Spt ^R , Tmp ^R	Tn916-like	t034/CC398	EU918672
7611472-1	Lamb/2004	Pen ^R , Cip ^R	Tn916-like	t034/CC398	EU918687
9b	Pig/2007	Pen ^R , Str ^R , Ery ^R , Spt ^R , Tmp ^R , Cef ^R , Met ^R	Tn916-like	t034/CC398 ^a	EU918691
7215190-1	Pig/2000	Pen ^R , Str ^R , Tmp ^R	Tn916-like	t034/CC398	EU918674
7215311-1	Pig/2000	Pen ^R , Str ^R	Tn916-like	t034/CC398	EU918675
7311242-1	Pig/2001	Ery ^R , Str ^R , Spec ^R	Tn916-like	t034/CC398	EU918676
7312330-1	Pig/2001	Pen ^R , Str ^R , Spec ^R	Tn916-like	t034/CC398	EU918677
7412791-1	Pig/2002	Pen ^R , Str ^R , Tmp ^R	Tn916-like	t034/CC398	EU918678
7413093-4	Pig/2002	Pen ^R , Ery ^R , Tia ^R , Tmp ^R	Tn916-like	t034/CC398	EU918679
7413532-2	Pig/2002	Pen ^R	Tn916-like	t011/CC398	EU918680
7413714-1	Pig/2002	Pen ^R , Tmp ^R	Tn916-like	t034/CC398	EU918681
7413727-1	Pig/2002	Pen ^R , Ery ^R	Tn916-like	t034/CC398	EU918682
7414035-2	Pig/2002	Pen ^R , Str ^R , Ery ^R	Tn916-like	t034/CC398	EU918683
7512166-1	Pig/2003	Pen ^R , Str ^R , Ery ^R , Tmp ^R	Tn916-like	t034/CC398	EU918684
7512986-1	Pig/2003	Ery ^R , Spt ^R	Tn916-like	t034/CC398	EU918685
7611280-5	Pig/2004	Pen ^R , Tmp ^R	Tn916-like	t011/CC398	EU918686
7611995-1	Pig/2004	Pen ^R , Str ^R , Ery ^R , Spt ^R , Tia ^R , Tmp ^R	Tn916-like	t034/CC398	EU918688
7612628-4	Pig/2004	Ery ^R , Spt ^R	Tn916-like	t571/CC398	EU918689
7711730-1	Pig/2005	Pen ^R , Spt ^R , Tia ^R , Tmp ^R	Tn916-like	t034/CC398	EU918690

^a : Determined by MLST

^b : Isolates suspected to be from the same outbreak.

Pen^R: penicillin, Str^R: streptomycin, Gen^R: gentamicin, Ery^R: erythromycin, Met^R: methicilin, Cip^R: ciprofloxacin,
 Chl^R: chloramphenicol, Spt^R: spectinomycin, Sul^R: sulphamethoxazo, Tia^R: tiamulin, Tmp^R: trimethoprin, Cef^R: Cefitfur,
 Fus^R : fusidic acid

532

533

534 **Table 3.** Primers used in this study

Primer number/name	Sequence	Reference
266 (Tet(M)-1)	5'-GTTAAATAGTGTTCTTGGAG-3'	27
267 (Tet(M)-2)	5'-CTAAGATATGGCTCTAACAA-3'	27
327 (Tn916-1)	5'-GCCATGACCTATCTTATA-3'	21
328 (Tn916-2)	5'-CTAGATTGCGTCCAA-3'	29
1811 (intcw459-1)	5'-CCGATATTGAGCCTATTGATGTG-3'	This study
1812 (intcw459-2)	5'-GTCCATACGTTTCCTAAAGTCGTC-3'	This study
709 (TetM sekvens 6)	5'-TCGAGGTCCGTCTGAAC-3'	This study
804 (TET-Down-1)	5'-GTCGTCCAAATAGTCGGATA-3'	This study
323 (TetM-up)	5'-CTGGCAAACAGGTTC-3'	21
324 (TetM-down)	5'-TAGCTCATGTTGATGC-3'	This study
526 (tet M upstream)	5'-TTGAATGGAGGAAAATCAC-3'	21
525 (tet M sekvens-1)	5'-TACTTTCCCTAAGAAAGAAAGT-3'	21
540 (Tet M seq-3)	5'-GCAGAAATCAGTAGAATTGC-3'	21
307 (Revers TetM-2)	5'-TTGTTAGAGCCATATCTTAG-3'	21
1835 (F1F)	5'-CGTGCAAATCTAGGTTATG-3'	This study
1836 (F2F)	5'-CATGAAGGAGTGTAAGAATGA-3'	This study
1837 (F2R)	5'-GTGTCTTATACCATGGAAGGA-3'	This study
1838 (F3F)	5'-GAGCCTCTTTAATCGCT-3'	This study
1839 (F3R)	5'-CATATTCGTCTGTCATGC-3'	This study
1840 (F4)	5'-GCTAGTGCTTCCATTAAGGA-3'	This study

535

536

537

538


539

540


541

542

Figure 1



563 Figure 2



564

565

566

567

568

569

570

571


572

573

574

575

576




578

579

580

581

582



584

585

586

587

588

589

590

591

592

593

594 Figure

595 5:

